



Molecular Characterization and Phylogenetics in the *Rhizophora* Species Complex at Pichavaram, Tamil Nadu: A Potential Resource for Phyto-Pharmacology

Utchimahali M¹, Shanthi A¹, Maheswari P¹, Nithishkumar K¹, Haritha S¹ and Nagarajan B^{1*} 

¹Institute of Forest Genetics and Tree Breeding, (Indian Council of Forestry Research and Education), P.B. No. 1061, Forest Campus R.S. Puram P.O., Coimbatore - 641 002, India.

Abstract: Mangroves are spatially limited bioresources known for their potential pharmacological uses. The genus *Rhizophora* is a conspicuous one reported to have enormous biopotentials in treating inflammation, diabetes, and rheumatism. However, mangroves are recently experiencing severe threats due to the over-exploitation of valuable tannin resources, climate change, and other anthropogenic pressures. *R × annamalayana*, a natural hybrid in Pichavaram, is rare in its distribution and a source of bioactive compounds with anticancer properties. The pure nature of the hybrid makes it more vulnerable to extinction. Therefore, domestication, conservation, and sustainable use of valuable resources are needed on a date. Poor genetic structure in mangroves is reported to be one of the reasons for extinction. Hence, understanding the population genetic structure of the genus is of paramount importance. Therefore, the present study aimed to understand the population genetic structure of the *Rhizophora* species complex in Pichavaram with the following objectives, to carry out the molecular characterization of the putative hybrid and to examine the phylogenetic relationship to its parental species using microsatellite markers. The study identified that putative hybrid *R. × annamalayana* (mean $H_E = 0.592$) has more significant variability than the putative parents *R. apiculata* (mean $H_E = 0.611$) and *R. mucronata* (mean $H_E = 0.667$). The negative inbreeding coefficient value or Wright's Index in *R. × annamalayana* (-0.190) suggests ample variation among the putative hybrids. The putative hybrid is genetically more proximal to *R. apiculata* than *R. mucronata*. Phylogenetic studies indicate that ten samples out of fifteen were clustered with *R. mucronata*, and the rest five grouped with *R. apiculata* indicating the varying paternal and maternal combinations.

Keywords: Inbreeding Coefficient, Nei's Distance, Phylogenetics, and Wright's Index.

*Corresponding Author

Nagarajan B , Institute of Forest Genetics and Tree Breeding, (Indian Council of Forestry Research and Education), P.B. No. 1061, Forest Campus R.S. Puram P.O., Coimbatore - 641 002, India.

Received On 26 September, 2022

Revised On 8 November, 2022

Accepted On 22 November, 2022

Published On 2 January, 2023

Funding

This research did not receive any specific grant from any funding agencies in the public, commercial or not for profit sectors.

Citation

Utchimahali M, Shanthi A, Maheswari P, Nithishkumar K, Haritha S and Nagarajan B , Molecular Characterization and Phylogenetics in the *Rhizophora* Species Complex at Pichavaram, Tamil Nadu: A Potential Resource for Phyto-Pharmacology.(2023).Int. J. Life Sci. Pharma Res.13(1), L202-208 <http://dx.doi.org/10.22376/ijlpr.2023.13.1.L202-208>



I. INTRODUCTION

Mangroves are ecologically essential bioresources with pharmacologically valuable flora distributed in the intertidal regions of the tropical and subtropical zone (30°N – 30°E). This group of plants is facultative halophytes, adapted to the muddy, anaerobic environment, extreme temperatures, and tidal regimes¹. These physiological adaptive mechanisms allow these plants to metabolize various active compounds that provide resilience in adverse environments². These active compounds play a significant role in the treatment of human illness. Most mangrove plants are widely used as traditional therapeutics³. In addition, mangroves provide tangible products to millions of life systems in and around tropical countries³. Mangroves are also known for their numerous ecosystem services, carbon sequestration, and coastal protection^{4,5}. Globally, eighty-one mangrove species are recorded under thirty genera and seventy families⁶. *Rhizophora* is a frequently noticeable genus in most of the mangrove realm with enormous biopotentials to treat inflammation, diabetes, and rheumatism⁷. In India, the genus *Rhizophora* is represented by three species (*R. apiculata*, *R. mucronata*, and *R. stylosa*) and two putative hybrids (*R. × annamalayana* and *R. × lamarckii*). Earlier investigations demonstrated the Anti-HIV activities of *R. apiculata*, the anti-inflammatory and antimicrobial properties of *R. mucronata*, and the hybrid *Rhizophora × annamalayana*^{8, 9}. Despite this, the genus *Rhizophora* is reported to produce high levels of tannin that are a potential source for free radical scavenging activity^{10, 11}. The putative hybrid *Rhizophora × annamalayana* occurs in Pichavaram (Tamil Nadu), a deltaic mangrove habitat from the Indian East Coast¹². The hybrid nature and parentage of this species were initially described by Kathiresan¹³, based on its intermediate morphology to the putative parents¹⁴. Later several investigations confirmed the hybridity of the species using molecular and DNA-based tools¹⁵⁻¹⁹. The hybrids are gigantic, with more excellent growth rates than their parental species^{12,20}, which promises a sustainable source of bioactive metabolites. However, the hybrid *R. × annamalayana* is rare within Pichavaram. New regeneration of the said hybrid is extremely rare²¹. The taxon is male-sterile; however, known

to produce propagules (very rarely) that do not grow into individual trees²². The sterile nature of the hybrids makes them more vulnerable¹⁵. The study location Pichavaram is a RAMSAR site. Recently, it has been subjected to several climate change extremities (Thane - 2011, Vardah - 2016, Okchi – 2017, Gaja - 2018, etc.). The infrequent population of *R. × annamalayana* in this location desires conservation and domestication for the sustainable extraction of biopotential supplies. Adequate protection in any mangrove resources requires a complete understanding of genetic diversity, as it is a fundamental component to accessing biodiversity²³. Knowledge of genetic diversity and patterns in threatened populations is critical to determining their health status, survival into posterity, and long-term endurance²⁴. Both natural and anthropogenic factors generally determine the genetic diversity pattern among the population of taxa. Therefore, it is a prerequisite to generate empirical data on these patterns. Similarly, lesser considerations on this aspect could also hamper the conservation effects¹⁷. Thus, the present study aimed to understand the population genetic structure of the *Rhizophora* species complex in Pichavaram with the following objectives, to carry out the molecular characterization of the putative hybrid and to examine the phylogenetic relationship to its parental species using microsatellite markers.

2. MATERIALS AND METHODS

2.1 Study Site and Target Taxa

Field surveys were conducted at the study site Pichavaram (11°17'– 11°30'N; 79°45'–79°50'E), a deltaic mangrove habitat of the Indian East coast. Plant collections were made in all three *Rhizophora* species. Specimens were initially identified using the flora. Three specimens in each taxa were sampled and deposited in Fisher's Herbarium at the Institute of Forest Genetics and Tree Breeding, Coimbatore (Table 1). This study sampled fifteen *Rhizophora × annamalayana* accessions and three accessions in each parental species (Table 2).

Table 1. Details of Accessions submitted to the Fisher's Herbarium, Forest campus, Coimbatore, TN.

S. No	Species	Herbarium ID	Geo-specifics
1	<i>R. apiculata</i> Blume.	FRC 25050	11°25'73.0" N; 79°47'62.2" E
2		FRC 25051	11°26'39.2" N; 79°47'55.8" E
4	<i>R. mucronata</i> Poir.	FRC 25048	11°25'55.9" N; 79°47'36.4" E
5		FRC 25049	11°27'47.0" N; 79°47'45.3" E
7		FRC 25052	11°25'55.1" N; 79°48'11.5" E
8	<i>R. × annamalayana</i> Kathiresan.	FRC 25053	11°25'45.6" N; 79°48'31.5" E
9		FRC 25054	11°25'33.5" N; 79°48'72.0" E

2.2 Genomic DNA Extraction

Rapidly expanding young leaves from the apical portion of the branch were sampled and dried in silica gel before DNA extraction. Arboreeasy™ DNA extraction kit (Institute of Forest Genetics and Tree Breeding) was used for extracting genomic DNA using the standard protocol. The purity of the DNA was quantified using a nanodrop spectrophotometer (nanodrop Inc., USA).

2.3 PCR Amplification and Microsatellite Genotyping

Fourteen microsatellite markers developed by Shinmura²⁵ were adopted in this investigation. Eurofins Genomics India Pvt

Ltd, Bangalore, India, synthesized the primers. Six of the most polymorphic and repetitive primers (Table 3) were shortlisted based on their banding pattern. A reaction mixture of 10 µl containing 1 µl of *Taq* buffer A, 0.3 µl of MgCl₂, 0.3 µl of 0.5mM dNTP mix, 0.5 µl of each of the primer, and 1 µl of template DNA (5-10 ng/µl) was subjected to the cyclic thermal amplification using BIO-RAD (BioRad Inc., USA) thermal cycler for 30 cycles. The amplicons were resolved on a 6% Polyacrylamide gel electrophoresis (vertical), and the gels were subjected to silver staining²⁶. Resolved gels were photographed using a DSLR camera (Nikon 300, Japan), and the products were scored manually.

3. STATISTICAL ANALYSIS

Co-dominant data was analyzed using GenAIEx add-in in Microsoft office excel (2019)²⁷ to calculate the number of different alleles (N.A.), Number of effective alleles (N.E.), Observed Heterozygosity (H.O.), Expected Heterozygosity

(HE), Polymorphic Information Content (PIC), allelic richness (R.S.), and Weight's Index (FST)). Phylogenetic analysis of the data sets was done using the Unweighted Pair Group Method with Arithmetic Means (UPGMA) by deploying the software DARwin (6.0.13)²⁸.

Table 2. Details of *Rhizophora* accession used for the Microsatellite analysis

S. No	Name of the taxon	Tree ID	GBH (cm)	Tree Height (m)	Geo-specifics
1		RA1	27.5	9.0	11°25'73.0" N; 79°47'62.2" E
2	<i>R. apiculata</i>	RA2	33.0	12.0	11°26'39.2" N; 79°47'55.8" E
3		RA3	30.5	10.0	11°27'4.70" N; 79°47'45.3" E
4		RM1	35.5	12.0	11°25'55.9" N; 79°47'36.4" E
5	<i>R. mucronata</i>	RM2	34.5	11.0	11°27'47.0" N; 79°47'45.3" E
6		RM3	35.5	12.0	11°26'46.3" N; 79°47'54.8" E
7		RXA 1	39.0	15.0	11°25'55.1" N; 79°48'11.5" E
8		RXA 2	48.0	12.0	11°25'45.6" N; 79°48'31.5" E
9		RXA 3	37.0	11.0	11°25'33.5" N; 79°48'72.0" E
10		RXA 4	37.5	13.0	11°25'31.8" N; 79°48'81.0" E
11		RXA 5	40.5	14.0	11°25'31.0" N; 79°48'90.0" E
12	<i>R. × annamalayana</i>	RXA 6	35.0	12.0	11°25'28.3" N; 79°48'10.5" E
13		RXA 7	37.5	10.0	11°25'27.2" N; 79°48'98.0" E
14		RXA 8	36.0	12.0	11°25'27.0" N; 79°48'99.0" E
15		RXA 9	35.5	10.0	11°25'26.4" N; 79°48'10.2" E
16		RXA 10	32.5	11.0	11°25'25.4" N; 79°48'10.2" E
17		RXA 11	37.5	11.0	11°25'24.0" N; 79°48'10.6" E
18		RXA 12	31.5	10.0	11°25'22.9" N; 79°48'11.3" E
19		RXA 13	32.0	10.0	11°25'22.9" N; 79°48'11.3" E
20		RXA 14	33.5	7.0	11°25'22.0" N; 79°48'11.5" E
21		RXA 15	36.0	12.0	11°25'55.1" N; 79°48'11.0" E

Table 3. Details of SSR Loci deployed for genotyping studies in *Rhizophora* species complex in Pichavaram, TN.

NCBI accession Number ²⁶	SSR code	Sequence	SSR motif	Tm (C)
AB721972	RM 107	F GGTTTTCCCAGTCACGACGAACAAGCATGGCAGGTAAC R GTTTGCCCATTTGGAATATGTGT	(CT) ₁₃	54
AB721976	RM 111	F GGTTTTCCCAGTCACGACGAACCGTTACTCGCGTATGCT R GTTTCATTGCCTCCATTCCATT	(T.C.) ₁₃	54
AB721977	RM 112	F GGTTTTCCCAGTCACGACGTTGAAGGTTGCGGTGAAAT R GTTTACATTCTTACCCCTGCGCACT	(AG) ₁₃	54
AB721979	RM 114	F GGTTTTCCCAGTCACGACGATTGGCATAGGCCTTGAATC R GTTTGTGGCTCAATTGTTGGCTA	(AT) ₁₃	54
AB721982	RM 121	F GGTTTTCCCAGTCACGACGTGGCCTATAGAGAAAGCGGA R GTTTCCTTCAATCCAAACAGC	(ATC) ₁₂	56

4. RESULTS

Three in each of the parental species (*R. apiculata* and *R. mucronata*) and fifteen putative hybrids (*Rhizophora × annamalayana*) samples were subjected to genotyping using fourteen microsatellite primers²⁵. Out of fourteen primers, six Loci were consistent and repetitive across the electrophoresis experiment. These primers were used as a diagnostic tool to understand the variation among the hybrids and their relationship with parental taxa. Microsatellite profile of the loci RM 107 is shown in Figure 1. Descriptive statistics of the *Rhizophora* species complex based on the six polymorphic microsatellite markers are presented in table 4. The average

number of alleles (N.A.) per locus was 2.0 – 3.3. The loci RM 111 and RM 107 had maximum and minimum alleles, respectively. The Observed Heterozygosity (H.O.) and the Expected Heterozygosity (HE) ranged from 0.444 – 0.867 and 0.473 – 0.647, respectively. The highest expected Heterozygosity (HE) is associated with the locus RM 114. The allelic richness (R.S.) range was 1.90 – 2.84. Polymorphic Information Content (PIC) varied from 0.49 – 0.74. The loci RM 112 and RM 116 recorded the highest PICs values of 0.71 and 0.74, respectively. In the present investigation, Weight Index (FST) values were 0.021 – 0.364 and were found statistically significant ($p < 0.05$).




Fig 1: SSR – Microsatellite profile of the loci RM 107 (1, 18 & 19. *R. apiculata*, 2, 20 & 21. *R. mucronata*, 3 – 17. *R. x annamalayana*).

Table 4. Descriptive statistics of different loci used in the study among the *Rhizophora* species complex at Pichavaram, TN.

Name of the Loci	<i>N_A</i>	<i>N_E</i>	<i>H_O</i>	<i>H_E</i>	<i>PIC</i>	<i>F_{SA}</i>
RM 107	3.33	2.394	0.489	0.550	0.66	0.164
RM 111	2.00	1.930	0.600	0.481	0.49	0.021
RM 112	2.33	1.906	0.844	0.473	0.71	0.337
RM 114	3.33	2.847	0.689	0.647	0.70	0.076
RM 116	2.66	2.593	0.444	0.592	0.74	0.203
RM 121	2.33	2.314	0.867	0.553	0.64	0.135
Mean	2.66	2.331	0.656	0.549	0.657	0.156

Table 4 catalogues different measures of genetic diversity among different loci used in the study. Several different alleles (*N.A.*), number of effective alleles (*N.E.*), Observed Heterozygosity (*H_O*), Expected Heterozygosity (*H_E*), Polymorphic Information Content (*PIC*), allelic richness (*R.S.*), and Weight's Index (*F_{ST}*) the higher the *H_E* and *PIC* indicate higher diversity.

Table 5 Summary of the genetic diversity at the species level among the *Rhizophora* species complex at Pichavaram, TN.

Name Species	<i>n</i>	<i>N_A</i>	<i>N_E</i>	<i>H_O</i>	<i>H_E</i>	<i>F_{IS}</i>
<i>R. apiculata</i>	3	2.167	2.100	0.611	0.509	-0.167
<i>R. mucronata</i>	3	2.500	2.300	0.667	0.546	-0.222
<i>R. x annamalayana</i>	15	3.333	2.593	0.689	0.592	-0.190

Table 5 catalogs different measures of genetic diversity among the *Rhizophora* species complex at Pichavaram, Number of individuals (*n*), Number of different alleles (*N.A.*), number of effective alleles (*N.E.*), Observed Heterozygosity (*H_O*), Expected Heterozygosity (*H_E*). Wright's Index or Inbreeding coefficient value (*F_{ST}*), the higher the *H_E* and *PIC* indicate higher diversity. The levels of genetic diversity noticed within the species complex among the taxa were found to vary (Table 5). The number of alleles in *R. apiculata*, *R. mucronata*, and *R. x*

annamalayana was 2.167, 2.500, and 3.333, respectively. However, the number of effective alleles or allelic richness was more or less similar across species. *R. x annamalayana* had the maximum mean observed heterozygosity (*H_E* = 0.689), followed by *R. mucronata* (*H_E* = 0.667) and *R. apiculata* (*H_E* = 0.611). The expected heterozygosity and Weight Index of *R. apiculata*, *R. mucronata*, and *R. x annamalayana* were 0.509, 0.546, 0.592, and -0.167, -0.222, and -0.190, respectively.

Table 6. Nei's Genetic distances observed between the *Rhizophora* species complex in Pichavaram, TN

	<i>R. apiculata</i>	<i>R. mucronata</i>	<i>R. x annamalayana</i>
<i>R. apiculata</i>	0.000		
<i>R. mucronata</i>	0.438	0.000	
<i>R. x annamalayana</i>	0.219	0.140	0.000

Nei's genetic distance computed using the allele frequency data among the three species of the *Rhizophora* species complex of Pichavaram is provided in table 6. The values describe the distance and relationship between the analyzed taxa. The phylogenetic relationship based on the UPMGA tree diagram reveals that the putative hybrid is an intermediary to the parental species (Fig 2). Nei's distance between *R. apiculata*

and *R. x annamalayana* was recorded to be 0.219, while the distance between *R. mucronata* and *R. x annamalayana* was 0.140. Further, the distance between the parental species (*R. apiculata* and *R. mucronata*) was 0.438. The parental species *R. apiculata* and *R. mucronata* are in the north and south of the dendrogram, and the putative hybrid genotypes fit in between. The hybrid accessions RxA-1 to RxA 10 clustered with *R.*

mucronata, whereas accessions RxA-11 to RxA-15 were grouped with *R. apiculata*. Nei's genetic distances (Table 6) observed among the species indicate that the putative hybrid species is genetically more proximal to *R. apiculata* than *R. mucronata*.

5. DISCUSSION

Mangroves are potential forest genetic resources with enormous phyto-pharmacological values. However, these valuable resources are severely threatened due to anthropogenic and natural pressures^{29,30}. The decrease in the efficacy of antibiotics owing to predisposition has accelerated demand for alternative drugs³¹. Hence, effective strategies to manage genetic diversity in Forest Genetic Resources gain paramount importance. Studies on biotically pollinated and abiotically seed dispersed taxa such as *Ceriops*, *Kandelia* and *Bruguiera* show low genetic variations^{32,33}. However, *Rhizophora*, an abiotically pollinated and dispersed genus, is of concern. Earlier studies in the Pichavaram indicate low levels of genetic variation within the population among species^{15,16}. This investigation reveals that the putative hybrid *R. × annamalayana* (mean $H_E = 0.592$) has more significant variability than its putative parents *R. apiculata* (mean $H_E = 0.611$) and *R. mucronata* (mean $H_E = 0.667$). The genetic diversity values noted among the Pichavaram hybrid populations are higher than that of *R. mucronata* (mean $H_E = 0.354$) and *R. stylosa* (mean $H_E = 0.321$) populations in Malaysia³⁴. The negative inbreeding coefficient value or Wright's Index in *R. × annamalayana* (- 0.190) suggests ample variation within the putative hybrids. This could be because of the varying mating systems among the parental species. The observed genetic

distances are consistent with the earlier study in site¹⁰ using the dominant marker system. Ten samples out of fifteen clustered with *R. mucronata*, and the rest five grouped with *R. apiculata*, indicating the varying paternal and maternal combinations. Further, investigations on Chloroplast DNA and Mitochondrial DNA on these lines may present more insights. The hybrid *R. × annamalayana* draws its lineage from both the putative parents. Both the putative parents of the hybrid share the common niche and overlap in the phenology leading to the natural hybridization^{19,35}. The taxon *R. apiculata* is reportedly threatened in Southeast Asia due to overexploitation of the tannin resources³⁶. Similarly, *R. apiculata* in the study location also had less population size. Notably, the loss of genetic diversity in one of the putative parents could implicate the process of natural hybridization. Hence, genetic diversity management during rehabilitation could amplify their rate of success. Further studies on this aspect with more markers and samples would give better insights into understanding the adaptive variations of the *Rhizophora* species complex in Pichavaram. This study demonstrates significant variations within the *R. × annamalayana* samples at an intra-population level. This suggests that the ecosystem has generated genetically, significantly varying recombinants. The putative hybrid population in this location needs further characterization and conservation at the forest genetic resource level. Conventional breeding technologies could be adapted to generate intraspecific hybrids with pedigrees, perhaps this could aid in improving the hybrid population and create an avenue for tree breeding research on developing wide tannin-yielding varieties if appropriate tree breeding and (or) improvement approaches are deployed.




Fig 2. Dendrogram showing the genetic relationship of the Hybrid *R. × annamalayana* with its parental species

6. CONCLUSION

The study indicates high genetic diversity among the *Rhizophora* species complex in Pichavaram. This suggests that

the ecosystem has generated genetically, significantly varying recombinants. The putative hybrid population in this location needs further characterization and conservation at the forest genetic resource level.

7. ACKNOWLEDGEMENTS

The authors thank the Indian Council of Forestry Research and Education (ICFRE), Dehra Dun, for the financial support and duly acknowledge Dr. R. Yasodha, Group Co-ordinator Research, Institute of Forest Genetics and Tree Breeding (IFGTB), Coimbatore, for the support and encouragements.

8. AUTHORS CONTRIBUTION STATEMENT

Dr. B. Nagarajan hypothesized the study and provided valuable

10. REFERENCES

1. Saenger P. Mangrove silviculture and restoration. In: *Mangrove ecology, silviculture and conservation*. Berlin: Springer; 2002. p. 229-70.

2. Nebula M, Harisankar HS, Chandramohanakumar N. Metabolites and bioactivities of *Rhizophoraceae* mangroves. *Nat Prod Bioprospect*. 2013 Oct;3(5):207-32. doi: 10.1007/s13659-013-0012-0.

3. Liquete C, Piroddi C, Drakou EG, Gurney L, Katsanevakis S, Charef A et al. Current status and future prospects for the assessment of marine and coastal ecosystem services: a systematic review. *PLOS ONE*. 2013 Jul 3;8(7):e67737. doi: 10.1371/journal.pone.0067737, PMID 23844080.

4. Kathiresan K. Mangrove forests of India. *Curr Sci*. 2018 Mar 10;114(5):976-81. doi: 10.18520/cs/v114/i05/976-981.

5. Murdiyarno D, Purbopuspito J, Kauffman JB, Warren MW, Sasmito SD, Donato DC et al. The potential of Indonesian mangrove forests for global climate change mitigation. *Nature Clim Change*. 2015 Dec;5(12):1089-92. doi: 10.1038/nclimate2734.

6. Saenger P, Ragavan P, Sheue C-R, López-Portillo J, Yong JWH, Mageswaran T. Mangrove biogeography of the Indo-Pacific. In: *Sabkha ecosystems*. Berlin: Springer; 2019. p. 379-400. doi: 10.1007/978-3-030-04417-6_23.

7. Kusuma S, Kumar PA, Boopalan K. Potent antimicrobial activity of *Rhizophora mucronata*. *J Ecobiotechnology*. 2012.

8. Premanathan M, Arakaki R, Izumi H, Kathiresan K, Nakano M, Yamamoto N et al. Antiviral properties of a mangrove plant, *Rhizophora apiculata* Blume, against human immunodeficiency virus. *Antiviral Res*. 1999 Dec 1;44(2):113-22. doi: 10.1016/s0166-3542(99)00058-3, PMID 10669261.

9. Alikunhi NM, Kandasamy K, Manoharan C, Subramanian M. Insulin-like antigen of mangrove leaves and its anti-diabetic activity in alloxan-induced diabetic rats. *Nat Prod Res*. 2012 Jun 1;26(12):1161-6. doi: 10.1080/14786419.2011.562205, PMID 22017188.

10. Rahim AA. Physico-chemical characterization of mangrove tannins as corrosion inhibitors [thesis]. Universiti Sains Malaysia; 2005 Dec.

11. Sulaiman S, Ibrahim D, Kassim J, Sheh-Hong L. Antimicrobial and antioxidant activities of condensed tannin from *Rhizophora apiculata* barks. *J Chem Pharm Res*. 2011;3(4):436-44.

12. Ragavan P, Jayaraj RS, Saxena A, Mohan PM, Ravichandran K. Taxonomical Identity of *Rhizophorax annamalayana* Kathir and *Rhizophorax lamarckii* Montrouz. (*Rhizophoraceae*) in the Andaman and Nicobar Islands, India. *Taiwania*. 2015 Oct 1;60(4).

13. Kathiresan K. *Rhizophora annamalayana*: a new species of mangroves. *Environ Ecol*. 1995;13:240.

14. Jayatissa LP, Dahdouh-Guebas F, Koedam N. A review of the floral composition and distribution of mangroves in Sri Lanka. *Bot J Linn Soc*. 2002 Jan 1;138(1):29-43. doi: 10.1046/j.1095-8339.2002.00002.x.

15. Parani M, Rao CS, Mathan N, Anuratha CS, Narayanan KK, Parida A. Molecular phylogeny of mangroves III Parentage analysis of a *Rhizophora* hybrid using random amplified polymorphic DNA and restriction fragment length polymorphism markers. *Aquat Bot*. 1997 Sep 1;58(2):165-72. doi: 10.1016/S0304-3770(97)00003-X.

16. Lakshmi M, Parani M, Parida A. Molecular phylogeny of mangroves IX: Molecular. *Aquat Bot*. 2002 Nov 1;74(3):201-17. doi: 10.1016/S0304-3770(02)00105-5.

17. Lo EYY. Testing hybridization hypotheses and evaluating the evolutionary potential of hybrids in mangrove plant species. *J Evol Biol*. 2010 Oct;23(10):2249-61. doi: 10.1111/j.1420-9101.2010.02087.x, PMID 20796134.

18. Kumar BA, Thangaraj M, Kathiresan K. Cross-species amplification of microsatellite loci in three mangrove species of India. *Int J Adv Biotechnol Res*. 2011;2(2):240-3.

19. Sahu SK, Singh R, Kathiresan K. Deciphering the taxonomical controversies of *Rhizophora* hybrids using AFLP, plastid and nuclear markers. *Aquat Bot*. 2015 Aug 1;125:48-56. doi: 10.1016/j.aquabot.2015.05.002.

20. Tyagi A.P. Cytogenetics and reproductive biology of mangroves in *Rhizophoraceae*. *Aust J Bot*. 2002;50(5):601-5. doi: 10.1071/BT01080.

21. Kavitha S, Kathiresan K. Reproductive biology of the most-at-risk mangrove species (*Rhizophora annamalayana*) and its parental species. In: *Proceedings of the international conference: Meeting on Mangrove Ecology, Functioning and Management (MMM3)*; 2012 Jul. p. 2-6.

22. Muniyandi K, Natarajan R. Incidence of seedling formation in *Rhizophora lamarckii* Montr. at Pichavaram mangrove, Tamil Nadu, India. *J Bombay Nat Hist Soc*. 1985.

23. Zimmer M, Ajonina GN, Amir AA, Cragg SM, Crooks S, Dahdouh-Guebas F et al. When nature needs a helping hand: different levels of human intervention for mangrove (re-)establishment. *Front For Glob Change*. 2022;5. doi: 10.3389/ffgc.2022.784322.

24. Azman A, Ng K.K., Ng CH, Lee CT, Tnah LH, Zakaria NF et al. Low genetic diversity indicating the threatened status of *Rhizophora apiculata* (*Rhizophoraceae*) in Malaysia: declined evolution meets habitat destruction. *Sci Rep*. 2020 Nov 5;10(1):19112. doi: 10.1038/s41598-020-76092-4, PMID 33154411.

inputs during the manuscript preparation. Dr. A. Shanthi, designed the primers and offered practical guidance during lab research and manuscript preparation. Mr. M. Utchimahali conducted the experiment, gathered data and drafted the manuscript. Ms. P. Maheshwari, Mr. K. Nithishkumar and S. Haritha supported in gathering wet lab data and analysis.

9. CONFLICT OF INTEREST

Conflict of interest declared none.

25. Shimura Y, Wee AKS, Takayama K, Meenakshisundaram SH, Asakawa T, Onrizal et al. Isolation and characterization of 14 microsatellite markers for *Rhizophora mucronata* (Rhizophoraceae) and their potential use in range-wide population studies. *Conservation Genet Resour.* 2012 Dec;4(4):951-4. doi: 10.1007/s12686-012-9681-y.

26. Zhang J, Wu YT, Guo WZ, Zhang TZ. Fast screening of microsatellite markers in cotton with PAGE/silver staining. *Acta Gossypii Sinica.* 2000;12(5):267-9.

27. Peakall R, Smouse PE. genalex 6: genetic analysis in Excel. Population genetic software for teaching and research. *Mol Ecol Notes.* 2006 Mar;6(1):288-95. doi: 10.1111/j.1471-8286.2005.01155.x.

28. Perrier X, Jacquemoud-Collet JP 2006. Darwin software <http://darwin.cirad.fr/darwin>.

29. Ellison JC, Zouh I. Vulnerability to climate change of mangroves: assessment from Cameroon, Central Africa. *Biology.* 2012 Nov 6;1(3):617-38. doi: 10.3390/biology1030617, PMID 24832511.

30. Friess DA, Adame MF, Adams JB, Lovelock CE. Mangrove forests under climate change in a 2°C world. *WIREs Climate Change.* 2022;13(4):e792. doi: 10.1002/wcc.792.

31. Smith P, Hiney MP, Samuels OB. Bacterial resistance to antimicrobial agents used in fish farming: a critical evaluation of method and meaning. *Annu Rev Fish Dis.* 1994 Jan 1;4:273-313. doi: 10.1016/0959-8030(94)90032-9.

32. Takeuchi T, Sugaya T, Kanazashi A, Yoshimaru H, Katsuta M. Genetic diversity of *Kandelia candel* and *Bruguiera gymnorhiza* in the Southwest Islands, Japan. *J For Res.* 2001 Aug 1;6(3):157-62. doi: 10.1007/BF02767087.

33. Ge X.J., Sun M. Population genetic structure of *Ceriops tagal* (Rhizophoraceae) in Thailand and China. *Wetlands Ecol Manag.* 2001 Jun;9(3):213-9. doi: 10.1023/A:1011156707160.

34. Yan YB, Duke NC, Sun M. Comparative analysis of the pattern of population genetic diversity in three Indo-west Pacific *Rhizophora* mangrove species. *Front Plant Sci.* 2016 Sep 30;7:1434. doi: 10.3389/fpls.2016.01434, PMID 27746790.

35. Kathiresan K, Bingham BL. Biology of mangroves and mangrove ecosystems; 2001. p. 81-251.

36. Richards DR, Friess DA. Rates and drivers of mangrove deforestation in Southeast Asia, 2000-2012. *Proc Natl Acad Sci U S A.* 2016 Jan 12;113(2):344-9. doi: 10.1073/pnas.1510272113, PMID 26712025.